

State of Alaska  
Department of Fish and Game  
Nomination for Waters  
Important to Anadromous Fish

242-31-10120 Segment 17-01 Trib 17

AWC Volume SE SC SW W AR IN USGS Quad Seldovia B-4

Anadromous Water Catalog Number of Waterway 242-31-10120-2272-3007

Name of Waterway \_\_\_\_\_ USGS name \_\_\_\_\_ Local name \_\_\_\_\_

Addition X Deletion \_\_\_\_\_ Correction \_\_\_\_\_ Backup Information \_\_\_\_\_

For Office Use

Nomination # <u>94 264</u>	<u>[Signature]</u> Regional Supervisor	<u>1/19/94</u> Date
Revision Year: <u>'94</u>	<u>Ed Weim</u>	<u>12/27/93</u>
Revision to: Atlas _____ Catalog _____	<u>2. Grove</u>	<u>2/2/94</u>
Both <u>X</u>	Drafted	Date
Revision Code: <u>A-2d</u>		

OBSERVATION INFORMATION

Species	Date(s) Observed	Spawning	Rearing	Migration	Anadromous
<u>Pink Salmon - Adult</u>	<u>9-11-93</u>	<u>4</u>			<u>✓</u>
<u>Coho Salmon - Juvenile</u>	<u>9-11-93</u>		<u>300 est.</u>		<u>✓</u>
<u>Dolly Varden - Juvenile</u>	<u>9-11-93</u>			<u>4</u>	<u>✓</u>

**IMPORTANT:** Provide all supporting documentation that this water body is important for the spawning, rearing or migration of anadromous fish, including: number of fish and life stages observed; sampling methods, sampling duration and area sampled; copies of field notes; etc. Attach a copy of a map showing location of mouth and observed upper extent of each species, as well as any other information such as: specific stream reaches observed as spawning or rearing habitat; locations, types, and heights of any barriers; etc.

Comments: Coho were observed throughout the system up to the barrier which is a spring. Pink Salmon were observed to a point shown on the sketch. Stream width is one meter at the mouth and upper extent. With the exception of the beginning and the end of the stream, the average width is 1.75 to 2 meters. Gradient is 2 percent. Good spawning and rearing. Predominant stream substrate is gravel.

ALASKA DEPT. OF  
FISH & GAME

Name of Observer (please print) JEFF BARNHART

NOV 03 1993

Date: 10-13-93

Signature: Jeff Barnhart

REGION II

Address: 333 Raspberry Road

WATER AND RESTORATION  
DIVISION

Anchorage AK

This certifies that in my best professional judgement and belief the above information is evidence that this waterbody should be included in or deleted from the Catalog of Waters Important for Spawning, Rearing or Migration of Anadromous Fishes per AS 16.05.870.

Signature of Area Biologist: \_\_\_\_\_

Rev. 7/93

# STREAM HABITAT ASSESSMENT 1993 - SEGMENTS

STREAM: Rocky 15 SEGMENT: 15-02 <sup>17-01</sup> DATE: 7/11/93 TEAM: TD/WG  
 ANADROMOUS: ☒ n WIDTH (m): 1 - 1 LENGTH (m): \_\_\_\_\_ GPS DATE: 7/12/93 DIGITIZE: y n  
 WATERBODY: mainstem tributary lake/pond wetland Intertidal other: \_\_\_\_\_

FISH					WILDLIFE		
SPECIES	STAGE (A J U)	COUNT	METHOD (E V D)	COMMENTS	SPECIES	COUNT	COMMENTS
pink	A	4	V		Bull moose	1	
lake	J	500	V		cow moose	1	
coho	J	3	D				
b.v.	J	4	D	2-3"			

GRADIENT(%): 1 CHANNEL PROFILE: V C D E F  
 CHANNEL PATTERN: single multi braided  
 STREAM SUBSTRATE: (rank three most predominant types) BEDROCK \_\_\_\_\_ BOULDER \_\_\_\_\_ RUBBLE \_\_\_\_\_ COBBLE \_\_\_\_\_  
 GRAVEL 1 SAND \_\_\_\_\_ MUD/SILT 2 ORGANICS 3 OTHER: \_\_\_\_\_  
 STREAM COVER TYPE: ORGANIC DEBRIS X DEAD BRANCHES/TWIGS \_\_\_\_\_ LOGS X BOULDERS \_\_\_\_\_  
 CUT BANK X OVERHANGING VEGET. X OTHER: \_\_\_\_\_  
 STREAM COVER ABUNDANCE: none low medium high

RIPARIAN VEGETATION (three most abundant plants in order of dominance) within 20m of the banks:  
 OVERSTORY: Sitka spruce  
 UNDERSTORY: Salmonberry Rose willow  
 CANOPY ABOVE STREAM: none low medium high  
 GROWTH: mature secondary shrubs meadow muskeg Intertidal

TOTAL BARRIER? ☒ n BARRIER TO SPECIES: coho/pinks adults juveniles  
 TYPE: fall slide beaverdam logjam spring substrate HEIGHT (m): \_\_\_\_\_ DIST. FROM UPPER EXTENT (m): 0

PHOTO ROLL(s): <u>JB05</u>		VIDEO TAPE(s): _____	
FRAME	DESCRIPTION	DATE	DESCRIPTION
<u>26</u>	<u>mud segment looking upstream</u>		

Substrate: Bedrock (solid) Boulder >1' Rubble 6-12" Cobble 2-6" Gravel .1-2" Sand <.1"  
 (Please enter comments on the other side) Comments

with the exception of the beginning & end. most of this  
stream averages 1.75 - 2 meters in width



242-31-18126

DO NOT ENTER

## STREAM HABITAT ASSESSMENT 1993 - STREAMS

STREAM: Rocky IS Trib 1-01 QUAD: SELDONIA B-4 STAGE: H M L  
 LANDOWNER: Chenega CAC Eyak Totillek Pt. Graham English Bay (circle one)  
 DATE(s): 9/10/93 UTM ZONE: 5  
 GPS FILES: 18092218F

SKETCH (indicate UTM zones, if not uniform throughout the stream)




PHOTO ROLL(s):

FRAME

DESCRIPTION

VIDEO TAPE(s):

DATE

(Please enter comments on the other side)


Excellent Naring Stream for Coho Salmon.  
500+ Coho fry observed along 21 schools of 3-5" Dolly  
Varden. 28 adult pinks observed. Moisture winds  
thick, three alders; high bush cranberries & eventually  
ends in a small upland pond. A Culvert has been  
placed in the stream for the old road which is  
adjacent to the upper extent.

One trail breaks off to the north & runs 100M  
where it ends due to the stream going underground.  
Coho fry to the end.

Ideal moose habitat, one bull observed  
signs of browsing on alders; numerous tracks  
in the stream bed.

10:37 tried GPS - Couldn't pick up satellite  
went from 1-4.





242-31-10120

tab 17 (17-pl)

SCALE 1:63360

Ps  
Cor



# MEMORANDUM

## State of Alaska

DEPARTMENT OF FISH & GAME

TO: Ed Weiss  
Habitat Biologist  
Region II  
Habitat and Restoration Division  
Department of Fish and Game

DATE: November 3, 1993

FILE NO.:

TELEPHONE NO.: 267-2295

SUBJECT: Anadromous Stream  
Nominations  
and Corrections  
Project R-51

FROM: Kathrin Sundet *KS*  
Habitat Biologist  
Region II  
Habitat and Restoration Division  
Department of Fish and Game

Attached are anadromous stream nominations and corrections to be included in the Anadromous Waters Catalog for 74 streams surveyed in the fall of 1993 on private lands held by the Port Graham, English Bay and Seldovia Native Corporations on the outer Kenai Peninsula.

Streams were surveyed by the Alaska Department of Fish and Game, Habitat and Restoration Division personnel, Kathrin Sundet, Jeff Barnhart, Dan Grey, and Wes Ghormley as part of Exxon Valdez Oil Spill Restoration project R-51 aka SHA (Stream Habitat Assessment).

Streams were surveyed on foot from the intertidal zone to the upper extent of anadromous fish distribution. Adult salmon and Dolly Varden were visually identified and enumerated. Juvenile salmon were visually identified in the stream, and then captured by electroshocking, dipnet, or minnow trap to confirm identification. Sampling was conducted periodically along the stream to determine the presence of juvenile salmon. No attempt was made to determine the rearing population sizes of juvenile salmon, or to determine the total escapement of adult salmon in a stream.

Stream data are on file at the Alaska Department of Fish and Game, Habitat and Restoration office, 333 Raspberry Road, Anchorage, Alaska.

cc: Lance Trasky  
Don McKay  
Mark Kuwada

ALASKA DEPT. OF  
FISH & GAME

NOV 03 1993

REGION II  
HABITAT AND RESTORATION  
DIVISION